Indicator of Sedation Need (IOSN)

ANXIETY QUESTIONNAIRE TO BE COMPLETED BY THE PATIENT

	Can you te	ell us how anxious yo	u get, if at all,	with your dental	visit?	
Please indicate by putting a 'X' in the appropriate box						
1. If you went to your Dentist for TREATMENT TOMORROW, how would you feel?						
	Not	Slightly	Fairly	Very Anxious □	Extremely	
	Anxious	Anxious	Anxious	Anxious [_]	Anxious	
2. If you we	2. If you were sitting in the WAITING ROOM (waiting for treatment), how would you feel?					
	Not	Slightly	Fairly	Very	Extremely	
	Anxious 🗌	Anxious	Anxious	Anxious	Anxious	
3. If you we	3. If you were about to have a TOOTH DRILLED, how would you feel?					
	Not Anxious 🗌	Slightly Anxious	Fairly Anxious 🗌	Very Anxious □	Extremely Anxious	
	Anatous	Anxious	Anatous	Anatous	Anatous	
4. If you we	. If you were about to have your TEETH SCALED AND POLISHED, how would you feel?					
	Not	Slightly	Fairly	Very	Extremely	
	Anxious 🗌	Anxious [_]	Anxious [_]	Anxious [_]	Anxious [
5. If you we would yo						
	Not .	Slightly	Fairly	Very	Extremely	
	Anxious	Anxious	Anxious 🗌	Anxious	Anxious 🗌	

Humphris GM, Morrison T and Lindsay SJE. The Modified Dental Anxiety Scale: Validation and United Kingdom Norms. Community Dental Health 1995; 12:143-150.

Dentist to score Anxiety Questionnaire

Each of the five answers is scored as follows:

Not anxious = 1 Slightly anxious = 2 Fairly anxious = 3 Very anxious = 4 Extremely anxious = 5 So the total Questionnaire Score is a sum of all five items (range 5 to 25)

Indicator of Sedation Need (IOSN)

MATRIX TO BE COMPLETED BY THE DENTIST

1. Anxiety Questionnaire (MDAS) Rank Score Questionannaire Score is converted to Rank Score MDAS 5-9 (minimal anxiety) MDAS 10-12 (moderate anxiety) MDAS 13-17 (high anxiety) 3

MDAS 18-25 (very high anxiety)

2. Medical & Behavioural Indicator Rank Score	Please circle one
No medical or behavioural indicators	1
Systemic disorders (not of severity to exclude sedation) that may be exacerbated by treatment Fainting attacks/ hypertension/ angina/ asthma/ epilepsy/ other (please state)	
Systemic disorders that compromise ability to cooperate Arthritis/parkinsonism/ multiple sclerosis/ other (please state)	2, 3, or 4
As a rule of thumb ASA II would generally be 2 or 3 and an ASA III would result in a grade of 4	
Gag reflex	

3 Treatment Complexity Rank Score Please circle one This guidance is not exhaustive - if in doubt about score then please score higher value ROUTINE - Scale, single rooted extraction of 1 or 2 teeth, small soft tissue biopsy, 1 single quadrant restorations, crown preparations or anterior endodontic treatment INTERMEDIATE - Scale and root planning, multi-rooted tooth extraction, surgical 2 extraction without bone removal, apicectomy anterior tooth, 2 quadrant restorative, posterior endodontic treatment 3 COMPLEX - Periodontal surgery, surgical extraction with bone removal, apicectomy posterior tooth, multiple quadrant restorative, multiple posterior endodontics HIGH COMPLEXITY - Any treatment considered more complex than above or are 4 multiples of the above

Total Rank Score Source Descriptor Sedation Need 3-4 Minimal need No 5-6 Moderate No 7-9 High need Yes 10-12 Very high need Yes

SEDATION NEED 1 + 2 + 3 scores