

Please answer these case scenarios so you are able to present them on the study day.

Please state further investigations and treatment you would carry out for each case.

### **CASE 1**

A 76-year-old gentleman with a history of cardiac problems presents at your surgery in the afternoon. He has had the extraction of the 46 carried out by your colleague earlier in the day. He is bleeding from the socket steadily and a large blood clot is visible.

### **CASE 2**

A 26-year-old lady presents three days following the extraction of the 16 complaining of fluid coming out of her nose on drinking. What do you look for and how do you treat initially and also definitively?

### **CASE 3**

You are extracting the 27. The tooth fractures and you are elevating the roots. The mesiobuccal root disappears. What is your plan of action?

### **CASE 4**

One week following the removal of a 48 a 33-year-old male returns for review at your surgery complaining of numbness of both his right lip and tongue. What do you do?

### **CASE 5**

You are extracting a 28 and hear a crack and see the palatal mucosa adjacent to the tooth tearing. What is happening? How do you proceed? What post operative instructions do you give to the patient?

### **CASE 6**

You are extracting a 55 on a 10-year-old child. As the tooth is extracted it slips out of the forceps and disappears. What do you do?

### **CASE 7**

You are having difficulty in extracting a 37 on a large 55-year-old male. You persist with elevating and forceps extraction until you hear a crack and the patient complains that his bite now feels wrong. What may have happened? How do you investigate and treat it? How may the problem have been avoided?

### **CASE 8**

A 73-year-old returns to your surgery following the removal of a 47 five days earlier. She complains she has a lot of pain, is unable to sleep and feels unwell. What clinical signs must you look for? What are the possible diagnoses? How would you treat the patient?

### **CASE 9**

A 22-year-old lady returns to your practice following a difficult surgical removal of a 38 five days post operatively. She complains that she cannot open her mouth. Her inter incisal opening is 5 mm. What may have caused this and how do you treat it?